

HUBUNGAN PEMBERIAN ASI EKSKLUSIF DENGAN FREKUENSI DIARE PADA ANAK USIA 6-24 BULAN DI DESA ARGOREJO SEDAYU BANTUL

Ahmad Antoni¹, Wahyuningsih², Sulistiyawati³.

Intisari

Latar Belakang Masalah : Diare merupakan salah satu penyakit utama pada bayi di Indonesia sampai saat ini dan menempati urutan ke tiga penyebab kematian bayi. Dilihat dari distribusi umur balita penderita diare di tahun 2010 didapatkan proporsi terbesar adalah kelompok umur 6-11 bulan yaitu sebesar 21,65%, lalu kelompok umur 12-17 bulan sebesar 14,43%. Salah satu penyebabnya adalah tidak diberikan ASI secara eksklusif selama 6 bulan. Persentase pola menyusui pada bayi umur 0 bulan adalah 39,8% menyusui eksklusif, 5,1% menyusui predominan, dan 55,1% menyusui parsial. Persentase menyusui eksklusif semakin menurun dengan meningkatnya kelompok umur bayi. Pada bayi yang berumur 5 bulan menyusui eksklusif hanya 15,3%, menyusui predominan 1,5% dan menyusui parsial 83,2%.

Tujuan penelitian : Untuk mengetahui hubungan pemberian ASI eksklusif dengan frekuensi diare pada anak usia 6-24 bulan di Desa Argorejo Sedayu Bantul.

Metode penelitian : Penelitian ini merupakan jenis penelitian kuantitatif. Dengan menggunakan rancangan *Cross Sectional*. Populasi dalam penelitian ini Populasi pada penelitian ini adalah seluruh anak yang berusia 6 sampai dengan 24 bulan di Desa Argorejo Sedayu Bantul. Diambil dengan *Tabel Krejcie* yang berjumlah 62 orang. Teknik pengambilan sampel dengan tehnik *Cluster Random Sampling*.

Hasil Penelitian : Penelitian ini dianalisa dan disimpulkan uji statistik *Chi-Square* di dapatkan nilai $p=0,001$ ($<0,05$) yang berarti nilai $p < \alpha$, H_0 = ditolak, H_a = diterima yang artinya ada hubungan antara pemberian ASI eksklusif dengan frekuensi diare pada anak usia 6-24 bulan di Desa Argorejo Sedayu Bantul.

Kesimpulan : Terdapat hubungan yang signifikan antara pemberian ASI eksklusif dengan frekuensi diare pada anak usia 6-24 bulan di Desa Argorejo Sedayu Bantul.

Kata Kunci: ASI eksklusif, Anak Usia 6-24 Bulan, Frekuensi Diare.

¹ Mahasiswa Program Studi Ners STIKES Alma Ata Yogyakarta

² Dosen Program Studi Ners STIKES Alma Ata Yogyakarta

³ Dosen Program Studi Ners STIKES Alma Ata Yogyakarta

THE PROVISION OF EXCLUSIVE BREASTFEEDING RELATIONSHIP WITH THE FREQUENCY OF DIARRHEA IN CHILDREN AGED 6-24 MONTHS IN THE VILLAGE ARGOREJO SEDAYU BANTUL

Ahmd Antoni¹, Wahyuningsih², Sulistiyawati³.

Abstract

Background problem: Diarrhea is one of major diseases in infants in Indonesia until now and ranked into three the cause of infant mortality. Viewed from the distribution of patients age toddler diarrhea in 2010 was obtained is the proportion of the largest age group 6-11 months namely 21,65 %, then 12-17 age group months worth 14,43 %. One cause was not given exclusively breast-fed for 6 months. The percentage of the pattern of breastfeeding in infants age 0 month is 39,8 % exclusive breastfeeding, 5.1 % breastfeeding predominant, and lactation 55.1 % partial. The percentage of exclusive breast-feeding skills with increasing age group baby. In infants who aged 5 months breastfeeding exclusively only 15.3 %, breastfeeding predominant 1.5 % and lactation partial 83,2 %.

Research objectives: To know the relationship the provision of breastfeeding exclusively with the frequency of diarrhea in children aged 6-24 months in the village argorejo sedayu bantul.

Methods: The research is quantitative the kind of research. Using cross sectional design. Population in this study population in this study were those with children ages six to 24 months in the village argorejo sedayu bantul. Taken with Krejcie table which consisted of 62 people. The sample collection technique technique with clusters of random sampling.

Results: This study were analysed and it can be concluded in the value of statistical tests chi-square, get the value of $p = 0.001$ (<0.05), which means the value of $p < \alpha$, H_0 = rejected, H_a = accepted that means that there is the relationship between the provision of breastfeeding exclusively with the frequency of diarrhea in children aged 6-24 months in the village argorejo sedayu bantul.

Conclusion: There are a significant relation between granting exclusive breastfeeding with the frequency of diarrhea in children aged 6-24 months in the village of argorejo sedayu bantul.

Keyword : Children aged 6-24 months, Exclusive breastfeeding, The frequency of diarrhea

¹ Student course of study STIKES Alma Ata Yogyakarta

² lecturers course of study STIKES Alma Ata Yogyakarta

³ lecturers course of study STIKES Alma Ata Yogyakarta